





## Stage 2 T1D

### Disease Modifying Therapy Clinical Trials (Multi-site)

 DRUG	<p><b>A Study of Baricitinib for the Delay of Stage 3 Type 1 Diabetes in At-Risk Children and Adults (BARICADE-DELAY)</b> AGES 1-36 STAGE 2 T1D (AT LEAST 2 AUTOANTIBODIES) ORAL PILL <b>NOT YET RECRUITING</b></p>	
---	---	---



## Stage 2 T1D


### Disease Modifying Therapy Clinical Trials (Single site)




 DRUG	<p><b>GLP-1Ra Impact on Metabolic Outcomes in Stage 2 T1DM While Receiving Teplizumab (GLP-TEP)</b> AGES 12-50 STAGE 2 T1D (AT LEAST 2 AUTOANTIBODIES) ORAL PILL VANDERBILT UNIVERSITY MEDICAL CENTER, NASHVILLE,TN</p>	
---	---	--

## Stage 3 T1D

### Disease Modifying Therapy Clinical Trials (Multi-site)

 DRUG	<p><b>A Study of Baricitinib to Preserve Beta Cell Function in Children and Adults Newly Diagnosed With Type 1 Diabetes (BARICADE-PRESERVE)</b> AGES 1-36 WITHIN 8 WEEKS OF DIAGNOSIS ORAL PILL <b>NOT YET RECRUITING</b></p>	
---	---	---

 DRUG	<p><b>A Study to Investigate Efficacy and Safety of Teplizumab Compared With Placebo in Participants 1 to 25 Years of Age With Stage 3 Type 1 Diabetes (βETA PRESERVE)</b></p> <p>AGES 1-25 WITHIN 8 WEEKS OF DIAGNOSIS IV INFUSION</p>	
 DRUG	<p><b>Targeting Type 1 Diabetes Using POLyamines (TADPOL)</b></p> <p>AGES 4-40 WITHIN 100 DAYS OF DIAGNOSIS ORAL PILL</p>	
 DRUG	<p><b>Precision Administration of Anti-thymocyte Globulin With or Without Verapamil</b></p> <p>AGES 6-36 WITHIN 100 DAYS OF DIAGNOSIS IV INFUSIONS AND ORAL PILL</p>	
 DRUG	<p><b>Rezpegaldesleukin in New Onset Type 1 Diabetes Mellitus</b></p> <p>AGES 8-45 WITHIN 100 DAYS OF DIAGNOSIS INJECTIONS <b>NOT YET RECRUITING</b></p>	

 DRUG	<p><b>A Randomized Phase 1/2 Trial of Low Dose Anti-thymocyte Globulin (ATG) With Subsequent Adalimumab or Verapamil in New Onset Type 1 Diabetes (WAVE T1D)</b>  AGES 9-21  WITHIN 6 MONTHS OF DIAGNOSIS  IV INFUSION  <b>NOT YET RECRUITING</b></p>	
 DRUG	<p><b>CNP-103 in Adolescent and Adult Subjects Ages 12-35 With Recently Diagnosed (Within 6 Months) Stage 3 Type 1 Diabetes (T1D)</b>  AGES 12-35  DIAGNOSED FOR LESS THAN 6 MONTHS  IV INFUSIONS</p>	
 DRUG	<p><b>A Phase III Study to Investigate if the Study Drug Diamyd Can Preserve Insulin Production and Improve Glycemic Control in Patients Newly Diagnosed With Type 1 Diabetes (DIAGNODE-3)</b>  AGES 12-28  WITHIN 6 MONTHS OF DIAGNOSIS  LYMPH NODE INJECTION</p>	
 DRUG	<p><b>FrexalimAB in Preservation of Endogenous insULIN Secretion Compared to Placebo in adULts and Adolescents on Top of inSulin Therapy (FABULINUS)</b>  AGES 12-35  WITHIN 3 MONTHS OF DIAGNOSIS  INFUSION FOLLOWED BY INJECTIONS</p>	




 DRUG	<p><b>Treatment of Type 1 Diabetes With Anti-OX40L Bispecific With Anti-TNF Activity In a Single Nanobody® Molecule (T1D OBTAIN)</b>  AGES 12-35  WITHIN 90 DAYS OF DIAGNOSIS  INJECTION</p>	
 DRUG	<p><b>SAFety and Efficacy of Human Anti-thymocyte ImmunoGlobULin SAB-142 ARresting Progression of Type 1 Diabetes (SAFEGUARD)</b>  PART A AGES 15-40  PART B AGES 5-40  WITHIN 100 DAYS OF DIAGNOSIS  IV INFUSION</p>	
 DRUG	<p><b>Denosumab for Type 1 Diabetes</b>  FEMALES AGES 18-50  MALES AGES 21-50  DIAGNOSED FOR LESS THAN 6 YEARS  INJECTION</p>	
 DRUG	<p><b>A Study of GNTI-122 in Adults Recently Diagnosed With T1D (POLARIS)</b>  AGES 18-45  WITHIN 120 DAYS OF DIAGNOSIS  SINGLE IV INJECTION</p>	

## Stage 3

### Disease Modifying Therapy Clinical Trials (Single site)

 DRUG	<p><b>PREBIOTICS IN PATIENTS WITH TYPE 1 DIABETES</b>          AGES 7+ (Lead Site)          AGES 7-17 (Subsites)          DIAGNOSED FOR LESS THAN 12 MONTHS          ORAL PILL          University of Calgary</p>	
 DRUG	<p><b>Cellular Therapy for Type 1 Diabetes Using Mesenchymal Stem Cells</b>          AGES 18-40          DIAGNOSED FOR LESS THAN 6 MONTHS          INFUSION          MEDICAL UNIVERSITY OF SOUTH CAROLINA</p>	
 DRUG	<p><b>An Adaptive Design of MTX228</b>          AGES 18-65          DIAGNOSED FOR AT LEAST 1 YEAR          ORAL PILL          UNIVERSITY OF ALBERTA</p>	
 DRUG	<p><b>MAS-1 Adjuvanted Antigen-specific Immunotherapeutic for Prevention and Treatment of Type 1 Diabetes</b>          AGES 18-45          DIAGNOSED FOR LESS THAN 2 YEARS          INJECTION          UNIVERSITY OF COLORADO, DENVER</p>	

## Trial Tools & Contact Information

 <b>TRIAL TOOLS</b>	<p><b>Clinical Trials Connection Tool</b> Match to a clinical trial in 60 seconds</p>	
 <b>CONTACT INFO</b>	<p><b>Clinical Trial Education Volunteers</b> Your key to participation! Connect with a local volunteer trained to answer any questions you may have.</p>	